

International Journal of Interdisciplinary and Multidisciplinary Research (IJIMR)

ISSN 2456-4567

Pathophysiological Outcomes of Ureteral Obstruction: An Update

Khalid Deshmukh¹, Gaaminepreet Singh¹

¹Department of Pharmacology, ISF College of pharmacy, Ghal Kalan, Ferozpur G.T. Road MOGA, Punjab

Corresponding author: **Gaaminepreet Singh**

Abstract

Renal impairment is frequently caused by blockage of the urine flow through urinary tract. Obstructive nephropathy can be induced by either unilateral or bilateral obstruction of ureter. For a long time, obstructive nephropathy was thought to be only a mechanical issue. Recent developments in cell and systems biology, have shown a complex physiopathology including a large number of molecular mediators of injury that contribute to cellular processes such as apoptotic cell death, cell injury, inflammation, and fibrosis. A chronic blockage causes tubular and glomerular obstruction. In addition to the acute impact on glomerular filtration rate and tubule function, interstitial damage also caused by the activation of several pathways. Chronic tubulointerstitial damage develops when tubulointerstitial injury progresses. Tubular atrophy, hypertrophy, inflammatory cell infiltration, and interstitial fibrosis are signs of renal injury. Based on current knowledge of the pathophysiology of renal damage, this minireview highlights the emerging events observed in obstructive nephropathy.

Keywords: 1 Ureteral obstruction, 2 Fibrosis, 3 Inflammation, 4 Apoptosis, 5 Tubular injury, 6 TGF- β

Introduction

Obstruction in flow of urine in any part of urinary system leads to obstructive nephropathy which can be induced by unilateral or bilateral ureteral obstruction. URO can be achieved by either reversible or irreversible ligation of ureter. Irreversible URO is characterised by ureter dilation, reduced renal blood flow, decrease in glomerular filtration rate, increasingly severe renal inflammation and interstitial fibrotic scarring while reversible URO (R-URO) is characterised by hydronephrosis, tubular dilation and inflammation[1].

In children, URO is a frequent cause of end-stage renal disease (ESRD), accounting for 21% of chronic kidney disease (CKD) and for 16% of children listed for transplantation. In adults, URO is a frequent cause of acute kidney injury (AKI). Up to 17% of AKI are secondary to URO [2].

Clinically, ureteral obstruction may cause by (a) urolithiasis, (b) congenital urinary tract malformations such as obstructive megaureter, uretero-pelvic junction obstruction, duplicated collecting system, horseshoe kidney, (c) immuno-rheumatologic diseases such as necrotizing vasculitis, periarteritis nodosa, kawasaki disease, henoch-schönlein purpura, eosinophilic ureteritis, (d) ureteral localization of infections like fungal infections, viral infections, bacterial infections, tubercular infections and (e) neoplastic intrinsic ureteral obstructions such as fibroepithelial polyps, ureteritis cystica, malignant neoplasms [3].

Irreversible ureteral obstruction

Many studies have reported that complete or irreversible ureteral obstruction is achieved by permanently obstructing the ureter by ligation with suture or by application of a ligating clip [1, 4-6] and is used to study renal inflammation and fibrosis [7]. It leads to blockage of urine flow thereby increasing hydrostatic pressure towards the kidney which triggers tubular cell death by renal fibrosis, interstitial inflammation, apoptosis and necrosis [8].

The sequence of events initiated rapidly in the kidney obstructed by complete UUO, leads within 24 h to reduced renal bloodflow and glomerular filtration rate followed by hydronephrosis, interstitial inflammatory infiltration (macrophages), and tubular cell death which leads to apoptosis and necrosis. Tubular epithelial cell death is caused by a number of stressors including ischemia, hypoxia and oxidant injury and axial strain caused by tubular dilatation [9].

Reversible ureteral obstruction

Reversible unilateral ureteral obstruction (R-UUO) is a prominent model of renal injury and used to study the structural and functional changes such as inflammatory and immune processes, cellular and tissue regeneration in kidneys after relief of the obstruction. R-UUO model is an advantageous model to assess kidney recovery after obstructive uropathy because it is a simple, easy, reproducible and more reliable method since it includes the nephrectomy of the contralateral kidney in order to invalidate its compensatory effect. [7].

In R-UUO model, the ureter is ligated for a suitable period of time sufficient to induce hydronephrosis, tubular dilation, inflammation and renal fibrosis. Then the ureteric obstruction is surgically reversed by anastomosis and kidney is allowed to decompress so that urinary flow gets restored toward the bladder. Due to this obstructed kidney which got hydronephrotic due to obstruction in urine flow becomes normal [1]. As a result, this model provides opportunity to investigators to explore the resolution of kidney inflammation together with key aspects of tissue repair [10].

Events involved in pathophysiological changes in UUO

A large number of studies has been performed to recognize the cellular and molecular effects obtained after ureteral obstruction [11]. Obstruction in ureter leads to changes in hydrostatic pressures, glomerular filtration, renal hemodynamics, and tubular function has been described well previously [12]. Apart from this various cellular events which include tubular dilation, phenotypic cellular transition, interstitial inflammation, followed by glomerulotubular injury and progressive interstitial fibrosis also found.

Tubular Injury

Tubular injury had occurred due to partial expansion in renal tubular lumen, degeneration of renal tubular epithelial cells which leads to tubular dilation [7]. Additionally, obstruction also rises hydrostatic pressure which gets transmitted back and eventually decreases the glomerular filtration rate (GFR) [12]. Moreover, obstructed kidney undergoes hydronephrotic with loss of brush border cells, tube formation with narrowing of renal parenchyma. In previous study in genetically modified β^6 -/- mice model, tubular injury was found in 5 days in wild type mice which gets proceeded in 14 days in renal cortex and medulla but significantly gets decreased in knockout mice [13].

Tubular Apoptosis

Apoptosis is a type of energy-dependent cell death that is brought on by harmful stimuli or a lack of survival factors. In UUO, tubular atrophy is caused by an increased rate of tubular cell death, with additional processes such as epithelial-mesenchymal transition playing a role (EMT) [14]. In UUO, apoptosis is observed in tubules as well as in interstitial cells in a time-dependent manner. Apoptosis in tubules is observed just a day after UUO induction whereas in interstitial cells it is found to be after 3 days [15]. Moreover, we found that Ang-II, TGF- β 1, members of the TNF cytokine superfamily,

oxidative stress, ATP depletion and mechanical stretch are also major triggerers of tubular apoptosis in UUO [16]

Tubular Fibrosis

Fibrosis is a well defined characteristic feature observed in ureteral obstruction. In fibrosis, there is a rise in numbers of activated fibroblasts and also various extracellular matrix components such as collagen I, III, and IV, proteoglycans, fibronectin gets accumulated [17]. It was reported that there is presence of α -SMA as well as expression of collagen-1 in tubulointerstitial space indicates the presence of interstitial myofibroblasts [7, 18]. Apart from that tubular fibrosis in obstructed nephropathy induced by UUO is also promoted by mitochondrial dysfunction and endoplasmic reticulum stress [13]. Fibrotic processes have been related to mitochondrial impairment also promotes kidney function deterioration in obstructive nephropathy [19-21]. Another study reported that breakdown of epithelial junctions and consequent loss of cell polarity are hallmarks of EMT [22].

Tubulointerstitial Inflammation

Inflammation is a complicated event that reflects local and/or systemic reactions to many stresses, and it typically allows disease resistance, tissue repair, and normal function restoration with the least amount of tissue damage feasible. Due to reversible obstruction in ureter, tubules get injured and initiate the secretion of macrophages, CD40, pro-inflammatory cytokines IL-2, MCP-1 and iNOS and also pro-fibrotic cytokine TGF- β 1 [7]. In previous study it was observed that interstitial inflammation due to infiltration of interstitial cells progressively enhanced from 12 hours after obstruction to up to 14 days [23,24]. A significant increase of adhesion molecules (ICAM-1 and VCAM-1), monocyte chemoattractive protein (MCP-1) and the glycoprotein osteopontin (OPN) is found after a few hours of ureteral blockage [23,25,26]. The increased expression of these molecules lasts for 7–10 days after the blockage has been removed. Within 5 days following blockage, ICAM-1 and OPN appear to have a function, although MCP-1 and VCAM-1 gene induction continues to rise [26-28].

Conclusion

In this review author try to convey about UUO and physiological and pathophysiological events occurs in tubules and kidney. Ureteral obstruction leads to tubular injury, apoptosis inflammation and fibrosis.

Consent for publication

The respective authors have declared their consent for the publication of this article.

Funding

The authors disclosed that no any funding or financial support was received for the publication of this article.

Declaration of Competing Interest

The authors declare that they have no potential conflicts of interest related to the contents of this article.

Acknowledgments

The authors are thankful for to Mr. Praveen Garg, Chairman and Prof. (Dr) G. D. Gupta, Director cum Principal, ISF College of Pharmacy, Moga, Punjab for his continuous support and encouragement.

References

- (1) Hesketh, E.E., Vernon, M.A., Ding, P., Clay, S., Borthwick, G., Conway, B. and Hughes, J., 2014. A murine model of irreversible and reversible unilateral ureteric obstruction. Journal of visualized experiments: JoVE, (94).

- (2) Trachtman, H., Weiser, A.C., Valderrama, E., Morgado, M. and Palmer, L.S., 2004. Prevention of renal fibrosis by spironolactone in mice with complete unilateral ureteral obstruction. *The Journal of urology*, 172(4 Part 2), pp.1590-1594.
- (3) Barros, A.N., Guiteras, R., Sola, A., Manonelles, A., Morote, J. and Cruzado, J.M., 2019. Reversal unilateral ureteral obstruction: A mice experimental model. *Nephron*, 142(2), pp.125-134. Da Silva, A.F., Silva, K., Reis, L.A., Teixeira, V.P. and Schor, N., 2015. Bone marrow-derived mesenchymal stem cells and their conditioned medium attenuate fibrosis in an irreversible model of unilateral ureteral obstruction. *Cell Transplantation*, 24(12), pp.2657-2666.
- (4) Liu, B., Ding, F.X., Liu, Y., Xiong, G., Lin, T., He, D.W., Zhang, Y.Y., Zhang, D.Y. and Wei, G.H., 2018. Human umbilical cord-derived mesenchymal stem cells conditioned medium attenuate interstitial fibrosis and stimulate the repair of tubular epithelial cells in an irreversible model of unilateral ureteral obstruction. *Nephrology*, 23(8), pp.728-736.
- (5) Ayyildiz, S.N., Ayyildiz, A., Benli, E., Cirrik, S., GERMİYANOĞLU, R.C., Noyan, T. and ÇIRAKOĞLU, A., 2017. The effect of trimetazidine on renal oxidative stress in partial and complete ureteral obstructions in a rat model. *Turkish Journal of Medical Sciences*, 47(3), pp.1012-1018.
- (6) Chevalier, R.L., Forbes, M.S. and Thornhill, B.A., 2009. Ureteral obstruction as a model of renal interstitial fibrosis and obstructive nephropathy. *Kidney international*, 75(11), pp.1145-1152. Bianchi, D., Vespasiani, G. and Bove, P., 2014. Acute kidney injury due to bilateral ureteral obstruction in children. *World journal of nephrology*, 3(4), p.182.
- (7) Hesketh, E.E., Vernon, M.A., Ding, P., Clay, S., Borthwick, G., Conway, B. and Hughes, J., 2014. A murine model of irreversible and reversible unilateral ureteric obstruction. *Journal of visualized experiments: JoVE*, (94).10.1038/ncpneph0098
- (8) CAPELOUTO, C.C. and SALTZMAN, B., 1993. The pathophysiology of ureteral obstruction. *Journal of endourology*, 7(2), pp.93-103.10.1002/biof.1673
- (9) Uceró, A.C., Gonçalves, S., Benito-Martin, A., Santamaría, B., Ramos, A.M., Berzal, S., Ruiz-Ortega, M., Egido, J. and Ortiz, A., 2010. Obstructive renal injury: from fluid mechanics to molecular cell biology. *Open access journal of urology*, 2, p.41. Gobe, G.C. and Axelsen, R.A., 1987. Genesis of renal tubular atrophy in experimental hydronephrosis in the rat. Role of apoptosis. *Laboratory investigation; a journal of technical methods and pathology*, 56(3), pp.273-281.
- (10) Misaki, T., Yamamoto, T., Suzuki, S., Fukasawa, H., Togawa, A., Ohashi, N., Suzuki, H., Fujigaki, Y., Oda, T., Uchida, C. and Kitagawa, K., 2009. Decrease in tumor necrosis factor- α receptor-associated death domain results from ubiquitin-dependent degradation in obstructive renal injury in rats. *The American journal of pathology*, 175(1), pp.74-83.
- (11) Jones, E.A., Shahed, A. and Shoskes, D.A., 2000. Modulation of apoptotic and inflammatory genes by bioflavonoids and angiotensin II inhibition in ureteral obstruction. *Urology*, 56(2), pp.346-351.
- (12) PIIS: Transforming Growth Factor--Dependent and -Independent Pathways of Induction of Tubulointerstitial Fibrosis in 6/Mice
- (13) Misseri, R., Meldrum, D.R., Dagher, P., Hile, K., Rink, R.C. and Meldrum, K.K., 2004. Unilateral ureteral obstruction induces renal tubular cell production of tumor necrosis factor- α independent of inflammatory cell infiltration. *The Journal of urology*, 172(4), pp.1595-1599.
- (14) Misseri, R., Meldrum, D.R., Dinarello, C.A., Dagher, P., Hile, K.L., Rink, R.C. and Meldrum, K.K., 2005. TNF- α mediates obstruction-induced renal tubular cell apoptosis and proapoptotic signaling. *American Journal of Physiology-Renal Physiology*, 288(2), pp.F406-F411.
- (15) Morrissey, J.J. and Klahr, S., 1997. Enalapril decreases nuclear factor κ B activation in the kidney with ureteral obstruction: Rapid Communication. *Kidney international*, 52(4), pp.926-933.
- (16) Lee, D.B., Huang, E. and Ward, H.J., 2006. Tight junction biology and kidney dysfunction. *American Journal of Physiology-Renal Physiology*, 290(1), pp.F20-F34.

- (17) Diamond, J.R., Kees-Folts, D., Ricardo, S.D., Pruznak, A. and Eufemio, M., 1995. Early and persistent up-regulated expression of renal cortical osteopontin in experimental hydronephrosis. *The American journal of pathology*, 146(6), p.1455.
- (18) Yoo, K.H., Thornhill, B.A., Forbes, M.S., Coleman, C.M., Marcinko, E.S., Liaw, L. and Chevalier, R.L., 2006. Osteopontin regulates renal apoptosis and interstitial fibrosis in neonatal chronic unilateral ureteral obstruction. *Kidney international*, 70(10), pp.1735-1741.
- (19) Ricardo, S.D., Levinson, M.E., DeJoseph, M.R. and Diamond, J.R., 1996. Expression of adhesion molecules in rat renal cortex during experimental hydronephrosis. *Kidney international*, 50(6), pp.2002-2010.
- (20) Morrissey, J.J. and Klahr, S., 1998. Differential effects of ACE and AT₁ receptor inhibition on chemoattractant and adhesion molecule synthesis. *American Journal of Physiology-Renal Physiology*, 274(3), pp.F580-F586.
- (21) Vielhauer, V., ANDERS, H.J., Mack, M., Cihak, J., Strutz, F., Stangassinger, M., Luckow, B., Gröne, H.J. and Schlöndorff, D., 2001. Obstructive nephropathy in the mouse: progressive fibrosis correlates with tubulointerstitial chemokine expression and accumulation of CC chemokine receptor 2-and 5-positive leukocytes. *Journal of the American Society of Nephrology*, 12(6), pp.1173-1187.
- (22) Weber, G.F., Ashkar, S., Glimcher, M.J. and Cantor, H., 1996. Receptor-ligand interaction between CD44 and osteopontin (Eta-1). *Science*, 271(5248), pp.509-512.